

Temporal correlator in YM_3^2 and reflection-positivity violation

A.R. Taurines*, A. Cucchieri† and T. Mendes†

**Instituto de Física Teórica, Universidade Estadual Paulista,
Rua Pamplona 145, São Paulo, SP, Brazil*

†*Instituto de Física de São Carlos, Universidade de São Paulo,
C.P. 369, 13560-970, São Carlos, SP, Brazil*

Abstract. We consider numerical data for the lattice Landau gluon propagator obtained at very large lattice volumes in three-dimensional pure $SU(2)$ Yang-Mills gauge theory (YM_3^2). We find that the temporal correlator $C(t)$ shows an oscillatory pattern and is negative for several values of t . This is an explicit violation of reflection positivity and can be related to gluon confinement. We also obtain a good fit for this quantity in the whole time interval using a sum of Stingl-like propagators.

INTRODUCTION

The reconstruction of a Gårding-Wightman quantum field theory from the corresponding Euclidean Green functions is possible only if they obey the Osterwalder-Schrader axioms. In particular, the requirement of positive definiteness of the norm in Hilbert space is expressed in Euclidean space by the *axiom of reflection positivity*. Introducing the *temporal correlator* of a 2-point function $D(p)$ as its partial Fourier transform

$$C(t) \equiv D(t, 0) = \int_{-\infty}^{\infty} \frac{dp_0}{2\pi} D(p) e^{ip_0 t}, \quad (1)$$

this axiom can be re-expressed simply as

$$C(t) > 0. \quad (2)$$

Thus, reflection positivity is violated if this temporal correlator assumes non-positive values.

The gluon propagator $D(p)$ is predicted to vanish at zero momentum in Landau gauge [1]. This implies that the real-space propagator $\tilde{D}(x-y)$ is positive and negative in equal measure, i.e. reflection positivity is maximally violated. This result can be interpreted as one *manifestation of confinement* in QCD, together with the Kugo-Ojima confinement criterion — which is equivalent to the horizon condition — and with the failure of the cluster decomposition (see [2] for a detailed discussion).

An explicit violation of reflection positivity for the Landau-gauge gluon propagator in YM_3^2 theory was presented in [3]. Here we review briefly the results obtained there and we try to fit the temporal correlator $C(t)$ in the whole t interval.




FIGURE 1. Temporal correlator $C(t)$ as a function of t for coupling $\beta = 6.0$ and lattice volume $V = 140^3$. The solid curve corresponds to the fit using the function reported in eq. (5).

VIOLATION OF REFLECTION POSITIVITY

On the lattice, the temporal correlator can be evaluated using

$$C(t) = \frac{1}{N} \sum_{k_0=0}^{N-1} e^{-2\pi i k_0 t / N} D(k_0, 0) , \quad (3)$$

where N is the number of points per lattice side and $D(k)$ is the propagator in momentum space. If the lattice action satisfies reflection positivity, it can be shown [4] that $C(t)$ is non-negative for all values of t . Furthermore, one can consider the quantity

$$G(t, a) = \frac{1}{a^2} [C(t)C(t+2a) - C(t+a)^2] , \quad (4)$$

where a is the lattice spacing. This function is also non-negative in the whole time interval if reflection positivity holds [3].

We consider the 3d $SU(2)$ Landau-gauge data presented in [5], obtained at very large lattice volumes ($V = 80^3, 140^3$) and for lattice couplings in the scaling region ($\beta = 4.2, 5.0, 6.0$). We find [3] that the temporal correlator $C(t)$ is negative for several values of t , showing a clear oscillatory behavior (see Fig. 1). We also checked that $G(t, 1)$ is negative for *all* values of t (as is the case of the Gribov-like propagator [3]). Thus, we find an *explicit violation of positivity* for the lattice Landau gluon propagator. Let us stress that this violation is clearly observable for the three lattice couplings and for the two lattice volumes considered.

TABLE 1. Parameters for the fitting formula (5), in lattice units.

	a_i	s_i	m_i	γ_i
i=1	-0.010	0.0013	-0.26	0.36
i=2	0.081	0.0081	-0.025	0.25
i=3	0.147	-0.013	-0.18	0.36

FITS FOR $C(T)$

The scaling behavior of the numerical data has been analyzed using the matching technique described in [5]. We find [3] that finite-size effects become important only for temporal separations $t \geq 3 fm$ and that all propagators become negative at $t \approx 0.7 fm$. In this scaling region the temporal correlator can be well fitted using a sum of two Gribov-like forms [3], with a light-mass scale $M \approx 325 MeV$.

Here we try to fit the data in the whole time interval using a sum of Stingl-like propagators [6]

$$D(p^2) = \sum_{i=1}^n \frac{a_i p^2 + s_i}{p^4 + m_i p^2 + \gamma_i^4}. \quad (5)$$

We find that a very good fit ($\chi/d.o.f. \sim 0.25$) is obtained considering $n = 3$. In Table 1 we report the fitting parameters for the lattice volume $V = 140^3$ with $\beta = 6.0$. The corresponding curve is shown in Fig. 1. Similar fits can be obtained in the other cases.

ACKNOWLEDGMENTS

This work was supported by FAPESP (Projects No. 00/05047-5 and 03/05259-0). Partial support from CNPq is also acknowledged (AC, TM).

REFERENCES

1. Zwanziger, D., *Nucl. Phys.*, **B364**, 127 (1991).
2. Alkofer, R., and von Smekal, L., *Phys. Rept.*, **353**, 281 (2001).
3. Cucchieri, A., Mendes, T., and Taurines, A. R., hep-lat/0406020 (2004).
4. Montvay, I., and Munster, G., *Quantum fields on a lattice*, Cambridge Univ. Press, Cambridge, 1994.
5. Cucchieri, A., Mendes, T., and Taurines, A. R., *Phys. Rev.*, **D67**, 091502 (2003).
6. Aiso, H., Fromm, J., Fukuda, M., Iwamiya, T., Nakamura, A., Stingl, M., and Yoshida, M., *Nucl. Phys. Proc. Suppl.*, **53**, 570 (1997).